

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification 7 :</b> <b>C12N 15/11, A61K 47/48, 48/00</b>	<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 00/32764</b> <b>(43) International Publication Date:</b> 8 June 2000 (08.06.00)
<b>(21) International Application Number:</b> PCT/EP99/08980 <b>(22) International Filing Date:</b> 22 November 1999 (22.11.99) <b>(30) Priority Data:</b> 98 403 015.5 2 December 1998 (02.12.98) EP <b>(71) Applicant (for all designated States except US):</b> I.D.M. IMMUNO-DESIGNED MOLECULES [FR/FR]; 172, rue de Charonne, F-75011 Paris (FR). <b>(72) Inventors; and</b> <b>(75) Inventors/Applicants (for US only):</b> MIDOUX, Patrick [FR/FR]; 21, rue du Poinçon, F-45100 Orléans (FR). PI-CHON, Chantal [FR/FR]; 1358, route d'Orléans, F-45640 Sandillon (FR). BELLO-ROUFAL, Mahajoub [FR/FR]; 5, rue de Tours, Résidence Les Hêtres, F-45072 Orléans Cedex 02 (FR). MONSIGNY, Michel [FR/FR]; 341, rue des Bouvreuils, F-45590 Saint-Cyr-en-Val (FR). <b>(74) Agents:</b> GROSSET-FOURNIER, Chantal et al.; Grosset-Fournier & Demachy, 20, rue de Maubeuge, F-75009 Paris (FR).		<b>(81) Designated States:</b> AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>With international search report.</i>
<b>(54) Title:</b> NEW OLIGOMERIC CONJUGATES LIABLE TO TRANSFER BIOLOGICAL MOLECULES INTO CELLS		
<b>(57) Abstract</b> <p>The invention relates to a positively charged oligomeric conjugate, containing an oligomer with a polymerization degree (PD) from 5 to 50, preferably 10 to 40 and more preferably 20, formed from monomeric components having free NH<sub>3</sub><sup>+</sup> in a number equal to or higher than 50 % of the polymerization degree. In particular, the invention provides new oligomeric conjugates of histidylated oligolysine liable to allow the transfer of oligonucleotides, peptides and oligosides into cells.</p>		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	R	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

## NEW OLIGOMERIC CONJUGATES LIABLE TO TRANSFER BIOLOGICAL MOLECULES INTO CELLS

---

5

The invention relates to new oligomeric conjugates liable to favor the transfer of biological molecules such as oligonucleotides, peptides and oligosides into cells.

10

The introduction of such molecules into cells is of great therapeutical interest.

Antisense oligonucleotides (ODN) and triplex forming oligonucleotides (TFO) are examples of attractive putative drugs in inhibiting or regulating gene expression in tumor cells and virus-infected cells.

15

Peptides from tumors and viruses are also attractive molecules in stimulating or eliciting a cell defense involving cytotoxic T lymphocytes against tumor and viruses after presentation by antigen presenting cells (APCs) such as dendritic cells, macrophages and B cells.

20

But to be effective these molecules must reach their target in the right intracellular compartments which are either the cytosol or the nucleus.

Until now, studies of the intracellular location of oligonucleotides point out that in the majority of cells, most of the oligonucleotides are confined inside vesicles once taken up and only a small amount succeeds in reaching their RNA and DNA targets in the cytosol and in the cell nucleus.

25

Because once in the cytosol, the oligonucleotides penetrate rapidly into the nucleus, the enhancement of the delivery of oligonucleotides into the cytosol upon cell uptake, is expected to increase their biological activity.

Oligonucleotide encapsulation into liposomes increases their delivery into the cytosol but efficiency is drastically reduced in the presence of serum.

30

Antigenic peptides can bind to MHC Class I molecules and be presented at the cell surface of APCs, upon their processing via proteasomes located in the cytosol.

Usually, peptides are taken up by APCs inside vesicles and delivered to lysosomes where they are either degraded or processed for an MHC Class II molecules presentation. Restricted MHC Class I molecules presentation by APCs requires that peptides must be introduced into the cytosol.

5 In French Patent n° 9613990 and PCT/FR97/02022, it has been shown that histidylated polylysine complexed with a nucleic acid is a system for cell transfection. The nucleic acid has a  $10^6$  to  $10^8$  of molecular weight. Polylysine is substituted at least 10% advantageously from 15% to 35% with molecules inducing membrane destabilization at acidic pH (mainly histidyl residues), and polylysine  
10 has a degree of polymerization of 15 to 900, particularly 200.

However, further studies have shown that the above-mentioned complex, which allows the transfection of cells by DNA, does not allow the transfer of an ODN (oligonucleotide).

One aim of the invention is to provide new positively charged oligomeric  
15 conjugates enabling the transmembrane passage of water soluble oligomers such as oligonucleotides, peptides and oligosides into the cytosol.

An other aim of the invention is to provide new oligomeric conjugates of substituted oligolysine liable to allow the transfer of oligonucleotides, peptides and oligosides into cells.

20 One advantage of the invention is that the formation of a complex between new oligomeric conjugates and oligoanions such as oligonucleotides, anionic peptides or anionic oligosides, is not required to allow their transfer into the cytosol and/or the cell nucleus.

Another advantage of the invention is that, although it is not excluded, the  
25 transfer of oligoanions such as oligonucleotides, anionic peptides, anionic oligosides (i.e. sulphated, phosphorylated, succinylated or sialylated oligosides) or a mixture thereof, does not require the formation of electrostatic complexes with the new positively charged oligomeric conjugates.

Another aim of the invention is to provide an *in vitro*, *ex vivo* and *in vivo*  
30 transfer process.

Another aim of the invention is to provide defined oligomeric compounds in which the nature of the monomeric compounds can be different from each other.

This has been achieved through the invention.

The invention, in one of its most general definitions, concerns a positively charged oligomeric conjugate containing an oligomer with a polymerization degree (PD) from 5 to 50, preferably 10 to 40 and more preferably 20, formed from monomeric components having free  $\text{NH}_3^+$  in a number equal to or higher than 50 % of the polymerization degree,

said oligomer being as follows :

- the free  $\text{NH}_3^+$  of the above-mentioned components are substituted in a ratio of at least 50 %, advantageously from 60 % to 95 %, particularly 80 to 90 % (this ratio being determined by nuclear magnetic resonance), by protonable residues in a weak acid medium, leading in such weak acid medium to a destabilization of cellular membranes,

- the above-mentioned protonable residues possess in addition the following properties :

- they contain a functional group enabling them to be linked to the above-mentioned oligomer,

- they do not correspond to a recognition signal recognized by a cellular membrane receptor,

- they can comprise at least one free  $\text{NH}_3^+$  group,

- the free  $\text{NH}_3^+$  of the above-mentioned monomers can be also substituted by an uncharged residue leading to a reduction of the number of positive charges in comparison to the same oligomeric conjugate, before substitution,

- molecules constituting a recognition signal recognized by a membrane cellular receptor may be present :

- either by substitution of some of the free  $\text{NH}_3^+$  of the above-mentioned monomers,

- either on some of the uncharged residues leading to a reduction of the number of charges,

- either on some of the above-mentioned protonable residues leading to a destabilization of the cellular membranes,

→ or by substitution of the free  $\text{NH}_3^+$  (if it is present) of the above-mentioned protonable residues leading to a destabilization of the cellular membrane,

provided that :

5           1) the total number of the non substituted  $\text{NH}_3^+$  is of at least 50 % of the polymerization degree,

2) the number of monomers initially carrying free  $\text{NH}_3^+$  is substituted in a ratio of at least 50 % of the polymerization degree by residues leading to a destabilization of the cellular membrane.

10           As it will result from the following, the first proviso corresponds to  $m \geq i/2$  and the second proviso corresponds to  $u \geq i/2$ .

Said oligomeric conjugate of the invention leads to the destabilization of the membranes and allows the transfer of the above-mentioned biomolecules in the cytosol and/or the nucleus of the cells.

15           According to a quite unexpected effect, the oligomeric conjugates of the invention allow the transfer of ODN into the cytosol and the nucleus, without allowing the transfection of cells by DNA.

The originality of the invention lies in the fact that the oligomer of the invention must be substituted to a level of more than 50 % with molecules inducing  
20 a membrane destabilization at acidic pH (amongst such molecules are histidyl residues) because oligomeric conjugates substituted at a level lower than 50 % are cytotoxic (see table 1, hereafter). Moreover, the toxicity of oligomeric conjugates substituted at a level lower than 50 % increases with the number of free  $\text{NH}_3^+$  of the unsubstituted lysyl residues.

25           For instance, for a oligomer of lysine of DP of 20 and carrying histidyl residues leading to the destabilization of the membranes, the above-said conditions can be expressed as follows :

- there must be at least 10 free  $\text{NH}_3^+$ , which come from the  $\alpha\text{-NH}_3^+$  and/or  $\epsilon\text{-NH}_3^+$ ,

30           - there are at least 10 histidyl residues, and even up to 20 histidyl residues carried by the side chain of the lysine residues which means, in the latter case, that

the minimum amount of free  $\text{NH}_3^+$  required, i.e. "at least" 10 free  $\text{NH}_3^+$  come only from the  $\alpha\text{-NH}_3^+$  functions of the histidyl residues.

The destabilization of membranes means a modification of membranes which leads either to the increase of their permeability with respect to low molecular weight (and possibly high molecular weight) molecules in solution, or the fusion with another membrane.

The membrane permeability can be measured as follows :

Cells are incubated at  $37^\circ\text{C}$  for 30 min in DMEM medium without serum in the presence of 0.5 mg/ml fluorescein-labelled dextran (Mw 4000) and in the absence or in the presence of an oligomeric conjugate. Cells are then washed and incubated for 30 min at  $37^\circ\text{C}$  in culture medium containing 10% serum. Cells are fixed for 5 min in PBS containing 4 % paraformaldehyde and the cell fluorescence localization is analysed under a fluorescent confocal microscope.

The fusion of membrane can be measured as follows :

~~The fusion of membrane can be measured by using liposomes according to~~  
Struck *et al.*, (Use of resonance energy transfer to membrane fusion. 1981 Biochemistry 20: 4093-4099).

Dioleoylphosphatidylcholine (DOPC) liposomes containing N-(7-nitrobenz-2-oxa-1,3-diazol-4-yl)-1,2-dihexadecanoyl-sn-glycero-3-phosphoethanolamine (NBD-PE) and octadecylrhodamine (R18) as fluorescent energy transfer donor and acceptor lipid probes, respectively, are mixed with non fluorescent liposomes and incubated in the absence or in the presence of oligomeric conjugates at various pH. Membrane fusion is evidenced by a decrease of the rhodamine fluorescence emission at 585 nm upon excitation at 470 nm, as a consequence of a decrease of the resonance energy transfer between NBD and rhodamine when the average spatial separation induced by membrane fusion increases.

The residues accounting for the destabilization of cellular membranes act through their property of being protonable in a weak acid medium.

The expression "weak acid medium" designates a medium the pH of which is lower than that of plasma or serum, i.e. a pH lower than 7.4.

Said medium can be either the extracellular medium or the lumen of intracellular compartments such as endosomes or lysosomes.

Said medium can be naturally acid or be acidified.

By way of example, the pH of said medium can be in the range of about 5 to about 7, in particular 5.5 to 6.5.

Transfer of the above-mentioned biomolecules into the cytosol and/or the cell nucleus, requires that both the oligomeric conjugate leading to the membrane destabilization and the above-mentioned biomolecules are present in the said medium.

In the present invention, the oligomeric conjugates and the above-mentioned biomolecules can be free or complexed.

Said positively charged oligomeric conjugate of the invention is liable to form a complex with at least one negatively charged oligoanion, the association between the oligoanion and the oligomeric conjugate being electrostatic in nature.

The expression according to which the protonable residues do not correspond to a recognition signal recognized by a cellular membrane receptor means that these residues are not used as ligands.

A molecule or a molecular complex is active as a recognition signal when it can be selectively recognized by a receptor, that is to say plays the role of a ligand, of an agonist, or of an antagonist. By recognition signal recognized by a cellular membrane receptor, one designates a ligand (molecule or molecular complexes) liable to be selectively recognized by said receptor.

According to an advantageous embodiment, the invention relates to a oligomeric conjugate, wherein the protonable residues leading to a destabilization of cellular membranes, present the additional properties :

- they are weak bases, the pK of which in aqueous medium is lower than 8, so that a proportion higher than 50 % of these bases linked to a cationic oligomer is not protonated at pH 7.4.

According to another advantageous embodiment, in the oligomeric conjugate of the invention, the protonable residues leading to a destabilization of cellular membranes, present the additional properties :

- they belong to the group of compounds comprising an imidazole ring,




- they belong to the group of quinolins,
- they belong to the group of pterins,
- they belong to the group of pyridins.

5

An example of quinolin is represented by the following formulae :

10




wherein  $R = (CH_2)_nCO_2H$ , in which  $n$  varies from 1 to 10, preferably from 1 to 3.

15

1 to 3.

Another example of quinolin is :

20




(3-quinoline carboxylic acid)

25



An example of pterin is represented by the following formula :

30



(pteroic acid)

Examples of pyridins are represented by the following formulae :



(nicotinic acid)

(quinolinic acid)

According to an advantageous embodiment, in the oligomeric conjugate of the invention, the protonable residues leading to a destabilization of the cellular membranes are :

- alkylimidazoles in which the alkyl radical comprises from 1 to 10, particularly from 2 to 6 carbon atoms, and in which only one of the nitrogen atoms of the imidazole ring is substituted.

According to an advantageous embodiment, in the oligomeric conjugate of the invention, the protonable residues leading to a destabilization of cellular membranes are chosen from :

histidine, 4-carboxymethyl-imidazole,

3-(1-methyl-imidazol-4yl)-alanine, 3-(3-methyl-imidazol-4yl)-alanine,


2-carboxy-imidazole, histamine, 3-imidazol-4yl)-L-lactic acid,

2-(1-methyl-imidazol-4yl)ethylamine, 2-(3-methyl-imidazol-4yl)ethylamine,

$\beta$ -alanyl-histidine-(carnosine), 7-chloro-4(amino-1-methylbutylamino)-quinoline,

N<sup>4</sup>-(7-chloro-4-quinoliny)-1,4-pentanediamine,  
 8-(4-amino-1-methylbutylamino)-6-methoxy-quinoline (primaquine),  
 N<sup>4</sup>-(6-methoxy-8-quinoliny)-1,4-pentanediamine, quininic acid,  
 quinoline carboxylic acid, pteric acid, nicotinic acid, quinolinic acid.

According to another advantageous embodiment, the oligomeric conjugate of the invention contains an oligomer of the following formula :



wherein



\* ai is an integer varying from 0 to 10,

\* bi is an integer varying from 0 to 10,


\* i = degree of polymerization from 5 to 50, and particularly 10 to 40, and preferably 20,

\* n = is an integer varying from 1 to 6, and preferably 4,


\* R represents in a ratio of 50 % to 100 % (corresponding to a number u)



or



or



m is an integer varying from 1 to 6,

n' is an integer varying from 0 to 6,

n'' is an integer varying from 0 to 6,

B is a weak base as defined above,

R' represents  $\text{NH}_3^+$  (corresponding to a number p),

or NH (corresponding to a number q) substituted by


-CO-CH<sub>3</sub>

-CO-(CHOH)rH

r being from 1 to 15 preferably  
2 to 7

-CO-(CH<sub>2</sub>)s-(CHOH)rH

r being from 1 to 15 preferably  
1 to 7 and s being from 1 to 6  
preferably 4



-SO<sub>2</sub>-Flu

-CO-Flu

-CS-NH-Flu

Flu being a fluorescent molecule

\* R represents in a ratio of 0 % to 50 % (corresponding to f : 0 < f ≤ u)


-  $\text{NH}_3^+$  (corresponding to a number j),

- NH (corresponding to a number k), substituted by

-CO-CH<sub>3</sub>



-CO-(CHOH)rH

r being from 1 to 15 preferably  
1 to 7



r being from 1 to 15 preferably  
1 to 7 and s being from 1 to 6  
preferably 4

5



Flu being a fluorescent molecule

10

- H (corresponding to a number h)


-  $(\text{CH}_2)_n\text{H}$ , n being an integer from 1 to 6 (corresponding  
to a number h)

-  $(\text{CH}_2)_n\text{OH}$  n being an integer from 1 to 6 (corresponding  
to a number h)

15

-  $(\text{CH}_2)_n\text{SA}'$

A' = H, CH<sub>3</sub> or



n being integer from 1 to 6 (corresponding to a number h)

20

with .  $i = u + j + k + h$

. total number of  $\alpha \text{NH}_3^+ = p = u - q$

. total number of  $\omega \text{NH}_3^+ = j = f - (k + h)$

. total number of  $\text{NH}_3^+ = m = p + j + 1$

25


with the proviso that :

1)  $u \geq i/2$

2)  $m \geq i/2$

30

According to another advantageous embodiment, the oligomeric conjugate of  
the invention contains an oligomer of the following formula :




wherein


\* i = degree of polymerization from 5 to 50, and particularly 10 to 40, and preferably 20,

\* n = is an integer varying from 1 to 6, and preferably 4,


\* R represents in a ratio of 50 % to 100 % (corresponding to u)



or



or



m is an integer varying from 1 to 6,

n' is an integer varying from 0 to 6,

n'' is an integer varying from 0 to 6,

B is a weak base as defined above,

R' represents  $\text{NH}_3^+$  (corresponding to a number p),

or NH (corresponding to a number q) substituted by

-CO-CH<sub>3</sub>

5

-CO-(CHOH)rH

r being from 1 to 15 preferably

1 to 7


-CO-(CH<sub>2</sub>)s-(CHOH)rH

r being from 1 to 15 preferably

1 to 7 and s being from 1 to 6

and preferably 4

10



-SO<sub>2</sub>-Flu

-CO-Flu

-CS-NH-Flu

15

Flu being a fluorescent molecule

\* R represents in a ratio of 0 % to 50 % (corresponding to f : 0 < f ≤ 1)

-  $\text{NH}_3^+$  (corresponding to a number j),

- NH (corresponding to a number k), substituted by

20

-CO-CH<sub>3</sub>

-CO-(CHOH)rH

r being from 1 to 15 preferably

1 to 7


-CO-(CH<sub>2</sub>)s-(CHOH)rH

r and s being from 1 to 15

preferably 1 to 7 and s being

from 1 to 6 and preferably 4

25



-SO<sub>2</sub>-Flu

-CO-Flu

30

-CS-NH-Flu

Flu being a fluorescent molecule

- H (corresponding to a number h)
- $(\text{CH}_2)_n\text{H}$ ; n being an integer from 1 to 6 (corresponding to a number h)
- 5        -  $(\text{CH}_2)_n\text{OH}$         n being an integer from 1 to 6 (corresponding to a number h)
- $(\text{CH}_2)_n\text{SA}'$          $\text{A}' = \text{H}, \text{CH}_3$  or  $\text{s} - \text{C}_5\text{H}_4\text{N}$

n being integer from 1 to 6 (corresponding to a number h)

10

- with
- $i = u + j + k + h$
  - total number of  $\alpha \text{NH}_3^+ = p = u - q$
  - total number of  $\omega \text{NH}_3^+ = j = f - (k + h)$
  - total number of  $\text{NH}_3^+ = m = p + j + 1$


15

with the proviso that :

- 1)  $u \geq i/2$
- 2)  $m \geq i/2$ .

20

According to another advantageous embodiment, the oligomeric conjugate of the invention contains an oligomer of the formula above defined, wherein



25

wherein




$$n' = n'' = 0$$

$$R' = \text{NH}_3^+$$

$$m = 1$$

$$B = \text{imidazole}$$

$$R = \text{NH-CO-CH-NH}_3^+$$



$$(f) R = \text{NH}_3^+$$


$$u = 12$$

$$j = 7$$

or

$$i = 19 \quad n = 4$$

$$(u) R = \text{NH-CO-(CH}_2\text{)}_{n'}\text{-CH-(CH}_2\text{)}_{n''}\text{-R'}$$



5

wherein


$$n' = n'' = 0$$

$$R' = \text{NH}_3^+$$

$$m = 1$$

$$B = \text{imidazole}$$

$$R = \text{NH-CO-CH-NH}_3^+$$




$$(f) R = \text{NH}_3^+$$

$$u = 16$$

$$j = 3$$

or

$$i = 19 \quad n = 4$$



wherein


$$n' = n'' = 0$$

$$\text{R}' = \text{NH}_3^+$$

$$m = 1$$

$$\text{B} = \text{imidazole}$$

$$\text{R} = \text{NH-CO-CH-NH}_3^+$$



$$(f) \text{ R} = \text{NH}_3^+$$


$$u = 19$$

$$j = 0$$

5

or

$$i = 19 \quad n = 4$$



wherein


$$n' = n'' = 0$$

$$\text{R}' = \text{NH}_3^+$$

$$m = 1$$

$$\text{B} = \text{imidazole}$$

$$\text{R} = \text{NH-CO-CH-NH}_3^+$$




$$(f) \text{ R} = \text{CO-CH}_3$$

$$u = 11$$

$$k = 8$$

or

$$i = 19 \quad n = 4$$



wherein



$$n' = n'' = 0$$

$$\text{R}' = \text{NH}_3^+$$

$$m = 1$$

$$\text{B} = \text{imidazole}$$

$$\text{R} = \text{NH-CO-CH-NH}_3^+$$




$$u = 15$$

$$k = 4$$

5

or

$$i = 19 \quad n = 4$$



wherein



$$n' = n'' = 0$$

$$\text{R}' = \text{NH}_3^+$$

$$m = 1$$

$$\text{B} = \text{imidazole}$$

$$\text{R} = \text{NH-CO-CH-NH}_3^+$$



$$r = 5$$



$$u = 12$$

$$k = 3$$

10

or

$$i = 19 \quad n = 4$$



5

wherein

$$n' = n'' = 0$$



$$m = 1$$


B = imidazole



$$u = 16$$

$$f = 4$$

$$k = 3$$




An example of mixed oligomer is the following :

10




15

20



wherein

$$n = 4$$



t varies from 1 to 6

As an example, when, in the above formula  $n = 4$ ,  $a_i = b_i = 0$ ,  $t = 1$ ,  
 5  $R' = \text{H}$ , the monomers are lysine and valine.

The invention also relates to a composition containing a mixture of at least one oligomeric conjugate as defined above, with at least one biological molecule, such as a peptide, an oligoside or an oligonucleotide derivative, or a mixture thereof.

10 In the composition of the invention, the oligomeric conjugates can be associated with a biological molecule, in particular an oligoanion, such as an oligonucleotide, an anionic peptide or an anionic oligoside, via electrostatic interactions.

15 An anionic oligoside can be a sulfated oligoside, succinylated oligoside, phosphorylated oligoside, sialylated oligoside or an oligoside containing pyruvylidenyl groups.

The invention also relates to a combined preparation containing as active substance the following individual components, in the form of a kit-of-parts :


- at least an oligomeric conjugate as defined above,
- 20 - at least one biological molecule, such as a peptide, an oligoside or an oligonucleotide, or a mixture thereof,

for the simultaneous, separate or sequential use, for the *in vitro*, the *in vivo* or the *ex vivo* transfer of said biological molecules into the cytosol and/or the cell nucleus.

The invention also relates to a complex between at least one oligoanion which can be an anionic peptide, an anionic oligoside or an oligonucleotide or a mixture thereof, a mixture of at least one non negatively charged biological molecule and of at least one oligoanion, and at least one positively charged oligomeric conjugate as defined above, the association between the oligoanion and the oligomeric conjugate being electrostatic in nature.


According to another advantageous embodiment, the biological molecule is chosen among oligonucleotides, peptides, oligosaccharides or a mixture thereof.

An oligonucleotide used in the invention can be of the following formula :



wherein  $i$  varies from 10 to 30, X represents O or S, B is a nucleic base U, A, T, G, C or a modified form such as a biotinyl or fluorescent labelled base which can be in  $\alpha$  or  $\beta$  anomeric position, R1 and R2 represent independently from

each other H, OH,  $(\text{CH}_2)_n\text{-A}$ ,  $[(\text{CH}_2)_2\text{-O}]_n\text{-CH}_2\text{-CH}_2\text{-A}$ , A being H, OH,  $\text{NH}_2$ ,  $\text{COOH}$ ,



, n being an integer from 1 to 6,

E represents H, OH,  $\text{OCH}_3$ ,  $\text{OCH}_2\text{CH}_3$ ,  $\text{O}(\text{CH}_2)_2\text{CH}_3$ ,  $\text{O}(\text{CH}_2)_3\text{CH}_3$ ,  $\text{O}(\text{CH}_2)_4\text{CH}_3$ ,  $\text{O-CH}_2\text{-CH}_2\text{-O-CH}_3$

As example of oligonucleotides, one may cite the following :

GEM 91

phosphorothioate ( $\text{X} = \text{S}$ ) oligonucleotide  $i = 25$

CTC TCG CAC CCA TCT CTC TCC TTC T

complementary to the AUG initiation site of gag HIV-1 gene

ISIS 1939

phosphorothioate ( $\text{X} = \text{S}$ ) oligonucleotide  $i = 19$


CCC CCA CCA CTT CCC CTC T

complementary to the 3' non coding region of ICAM-1 mRNA.


A mixture of an oligonucleotide and a peptide is defined as an oligonucleotide linked to a peptide, and a mixture of an oligonucleotide and an oligoside is defined as an oligonucleotide linked to an oligoside.

A mixture of an oligoside and a peptide is defined as an oligoside linked to a peptide.

A mixture of an oligonucleotide and a peptide or a mixture of an oligonucleotide and an oligoside used in the invention can have the following formulae :



wherein  $i$  varies from 10 to 30,  $X$  represents O or S,  $B$  is a nucleic base U, A, T, G, C or a modified form such as a biotinyl or fluorescent labelled base which can be in  $\alpha$  or  $\beta$  anomeric position,  $R_1$  and  $R_2$  represent independently from each other H, OH,  $(CH_2)_n-A$ ,  $[(CH_2)_2-O]_n-CH_2-CH_2-A$ ,  $A$  being H, OH,  $NH_2$ , COOH,



,  $n$  being an integer from 1 to 6,


or a peptide or an oligoside,

$E$  represents H, OH,  $OCH_3$ ,  $OCH_2CH_3$ ,  $O(CH_2)_2CH_3$ ,  $O(CH_2)_3CH_3$ ,  $O(CH_2)_4CH_3$ ,  $O-CH_2-CH_2-O-CH_3$




When R1 and/or R2 represent a peptide, the mixed oligoanion is a peptido-oligonucleotide, and when R1 and/or R2 represent an oligoside, the mixed oligoanion represents a glyco-oligonucleotide.

An example of oligonucleotide is a peptide nucleic acid (PNA) represented by the following formula :



wherein - R1 and R2 represent independently from each other H, OH, (CH<sub>2</sub>)<sub>n</sub>-A, [(CH<sub>2</sub>)<sub>2</sub>-O]<sub>n</sub>-CH<sub>2</sub>-CH<sub>2</sub>-A, A being H, OH, NH<sub>2</sub>, COOH,



, n being an integer from 1 to 6,

- i is an integer varying from 10 to 30.

5 In the composition of the invention, the oligonucleotide can be simple or double stranded, or can form a triplex (three strands) or a quadruplex (4 strands).

The invention also relates to the use of an oligomer conjugate as defined above, for the intracellular transfer of biological molecules into the cytosol or/and into the cell nucleus *in vitro*, *ex vivo* or *in vivo*.

10 The invention also relates to the use of an oligomeric conjugate as defined above or of a composition as defined above, or of a combined preparation as defined above, for the intracellular *in vitro*, *ex vivo* or *in vivo* transfer of a peptide, an oligoside or an oligonucleotide, or a mixture thereof, into the cytosol or/and in to the cell nucleus of cells.

15 The invention also relates to the use of an oligomeric conjugate as defined above or of a composition as defined above, or of a combined preparation as defined above, wherein the cells are chosen among muscular, epithelial, endothelial, or myeloid cells such as monocytes, macrophages, fibroblasts, leukocytes and granulocytes, osteoblasts, as well as dendritic, stem, neuronal, or  
20 dermal cells.

The invention also relates to a method for the *in vivo*, the *in vitro* or the *ex vivo* transfer of an oligonucleotide, wherein an oligonucleotide and an oligomeric conjugate as defined above, or of a composition as defined above, or of a combined preparation as defined above, are(is) contacted with a medium containing  
25 cells to be transferred, under conditions such that there is :

- transfer of an antisense oligonucleotide in the cytosol and/or the cell nucleus where it binds and blocks the complementary mRNA sequence,

- transfer of an oligonucleotide as activator into the cytosol where it depresses or activates a second messenger in the cytosol, or the corresponding gene  
30 in the nucleus,

- transfer into the cytosol and/or the cell nucleus of oligonucleotides corresponding to a repetitive bacterial type DNA sequence with stimulating or immunodepressive activity.

5 The transfer of an antisense oligonucleotide in the cytosol where it binds to the complementary mRNA sequence and blocks its traduction leading to inhibition of the synthesis of the gene product, can be carried out as described hereafter in the legends of Figures 1, 2 and 3.

10 The invention also relates to a method for the *in vivo*, the *in vitro* or the *ex vivo* transfer of an oligonucleotide, wherein an oligonucleotide and an oligomeric conjugate as defined above, or of a composition as defined above, or of a combined preparation as defined above, are(is) contacted with a medium containing cells to be transferred, under conditions such that there is :

15 - transfer into the cytosol and/or the cell nucleus of RNA or DNA oligonucleotide acting as decoys which inhibit gene expression by blocking the binding of regulatory factors to the authentic DNA region such as short RNA oligonucleotides corresponding to the HIV-TAR sequence inhibiting HIV expression and replication by blocking the binding of the HIV regulatory protein at Tat to the TAR region,

20 - transfer into the cytosol and/or the cell nucleus of ribozymes (RNA oligonucleotides) which inhibit gene expression by cleaving the mRNA.

The transfer into the nucleus of an oligonucleotide (triplex forming ODN, TFO) where it binds to target DNA at oligopurine sites where they form a triple-helical structure, leading to the inhibition of the gene expression, can be processed as hereafter described.

25 As an example of the specific TFO is an oligonucleotide 5'-A<sub>4</sub>GA<sub>4</sub>G<sub>6</sub>A-3' directed against the polypurine track (PPT) in the NEF-HIV-1 gene.

The transfer of the oligonucleotide as activator of the immune response can be carried out as follows :

30 As an example the double strand RNA polyinosinic-polycytidylic acid (poly(I:C)) for the increase of the tumoricidal activity of macrophages or for the stimulation of natural killer lymphocyte cytotoxicity.

The transfer of double stranded polynucleotide as activator of the immune response can be illustrated by the following :

Polyinosinic-polycytidylic acid (poly(I:C)) is known to increase the tumoricidal activity of macrophages *via* a TNF- $\alpha$  mediated cytolysis.

5 Thioglycolate-elicited peritoneal macrophages ( $2.5 \times 10^5$ ) are plated in 16 mm diameter well multiwell plates in serum free RPMI medium. Upon 2 h incubation at 37°C, non-adherent cells are discarded. Adherent cells are cultured in 0.5 ml of serum free RPMI medium in the absence or the presence of polyI:C and in the absence or the presence of histidylated oligolysine. Culture supernatants are  
10 collected after 24 h incubation at 37°C. Before testing, supernatants are rendered cell-free by centrifugation at 2000 g for 10 minutes. The cytotoxic activity of macrophage culture supernatants is determined by using L929 cells pretreated for 2 h at 37°C with 2  $\mu$ g/ml actinomycin D. Actinomycin D pretreated L929 cells are seeded in 96-wells microplates ( $4 \times 10^4$  cells in 0.05 ml serum free RPMI medium  
15 per well). After 3-4 h at 37°C, diluted supernatants from stimulated macrophages are added (0.05 ml) and the cells are incubated at 37°C for 18-20 h. Microplates are washed with PBS and the percentage of cell lysis is determined after staining the cells with 0.05 ml of crystal violet (0.2% in 2 % ethanol). The plates are washed with tap water and the dye is solubilized by adding 0.06 ml per well of 0.3  
20 % sodium dodecyl sulfate. Absorbance of each well is read at 570 nm. The % cytotoxicity is calculated according to  $(A_{NS} - A_S)/A_{NS} \times 100$  where  $A_{NS}$  and  $A_S$  are the absorbances of wells containing target cells incubated with supernatant dilutions of non stimulated and stimulated macrophages, respectively.

25 The invention also relates to a method for the *in vivo*, the *in vitro* or the *ex vivo* transfer of peptide, wherein a peptide and an oligomeric conjugate as defined above, or of a composition as defined above, or of a combined preparation as defined above, are(is) contacted with a medium containing cells to be transferred, under conditions such that there is a transfer of said peptide into the cytosol.

30 The transfer of a peptide can be illustrated by the following :

5 A) Cells are incubated for 4 h at 37°C with 1  $\mu$ M fluorescein-labelled peptide (F-S-CGEEDTSEKDEL) in the absence or in the presence of histidylated oligolysine. Cells are fixed with 2 % of p-formaldehyde, washed and mounted on slides in a PBS/glycerol mixture (1:1 v/v) containing 10 mg/ml DABCO (1,4 diazobicyclo-(2,2,2)-octane) as antifading agent. Cells are analyzed with a confocal microscope imaging system (MRC-1024, Bio-Rad) equipped with a Nikon Optiphot epifluorescence microscope.

10 B) Dendritic cells are incubated for 4 h at 37°C with 1  $\mu$ M c-myc epitope peptide (SMEQKLISEEDLNFELEA) in the absence or in the presence of histidylated oligolysine. Cells are fixed with 2 % of p-formaldehyde in the presence of 0.5 % saponine, washed and then incubated for 1 h with anti c-myc epitope monoclonal antibody (9E10) in PBS containing containing 10 mg/ml BSA and 0.1% saponin. Cells are washed and further incubated for 1 h in the presence of fluorescein-labelled anti-mouse IgG F(ab)' fragments in PBS containing containing 10 mg/ml BSA and 0.1% saponin. Cells are washed and mounted on slides in a PBS/glycerol mixture (1:1 v/v) containing 10 mg/ml DABCO (1,4 diazobicyclo-(2,2,2)-octane) as antifading agent. Cells are analyzed with a confocal microscope imaging system (MRC-1024, Bio-Rad) equipped with a Nikon Optiphot epifluorescence microscope.

The fixation of a peptide to intracellular cofactor can be carried as described in the following example:

25 An oligopeptide corresponding to a prostate specific antigen (PSA) epitope mixed to the oligomeric conjugate is transferred into the cytoplasm of macrophages.

30 The oligopeptide is fixed there to heat shock protein s (HSP90, HSP70) to form HSP-peptide complexes which are then re-expressed at the surface of macrophages. This complex formed with the HSP cofactor stimulate macrophages and enhance the immune response to the PSA antigen.

The invention also relates to a method for the *in vivo*, the *in vitro* or the *ex vivo* transfer of peptide, wherein a peptide and an oligomeric conjugate as defined

above, or a composition as defined above, or a combined preparation as defined above, in particular wherein the peptide is an antigenic peptide, are(is) contacted with a medium containing cells to be transferred, under conditions such that there is a transfer of said antigenic peptide in the cytosol of antigen presenting cells (macrophages, dendritic cells and B cells) where they are processed in proteosomes in order to bind to MHCI molecules, allowing the presentation of the antigenic epitope fixed on MHCI.

*In vitro* evaluation of antigen presentation can be illustrated by the following:

Dendritic cells were incubated for 4 h at 37°C with the nonadecapeptide (185-203) from the C-terminal part of the HIV-1 Nef protein containing the nonapeptide (190-198) (AFHHVAREL) in the absence or in the presence of an histidylated oligolysine. Cells are washed and further incubated for 24 h at 37°C in the absence of peptide and histidylated oligolysine. MHC class I presentation of peptide antigen was evaluated by Cr<sup>51</sup> cytotoxic assay by using a CTL clone sensible to the peptide. DCs were labelled with Cr<sup>51</sup> (target cells : T) and then incubated at 37°C for 4 h in the presence of the CTL clone (effector cells : E) at E/T ratios ranged from 1 to 100. The supernatants are collected and the radioactivity in the supernatant was recorded. The % of specific Cr<sup>51</sup> release is calculated according to  $(A_{NS} - A_S)/A_{NS} \times 100$  where  $A_{NS}$  and  $A_S$  are the radioactivity in supernatant dilutions of dendritic cells incubated in the absence and the presence of CTL cells, respectively.

The invention also relates to a method for the *in vivo*, the *in vitro* or the *ex vivo* transfer of a oligoside, wherein an oligoside and an oligomeric conjugate as defined above, or a composition as defined above, or a combined preparation as defined above, are(is) contacted with a medium containing cells to be transferred, under conditions such that there is a transfer of said oligoside into the cytosol and/or the cell nucleus.

An example of transfer of an oligoside can be illustrated by the following :

Cells are incubated for 4 h at 37°C with 0.5 mg/ml fluorescein-labelled dextrans (either Mw 4000 or Mw 70000) in the absence or in the presence of an oligomeric conjugate. Cells are washed with PBS, fixed with 2 % of p-formaldehyde, washed and mounted on slides in a PBS/glycerol mixture (1:1 v/v)

containing 10 mg/ml DABCO (1,4 diazobicyclo-(2,2,2)-octane) as antifading agent. Cells are analyzed with a confocal microscope imaging system (MRC-1024, Bio-Rad) equipped with a Nikon Optiphot epifluorescence microscope (Nikon, Tokyo, Japan) and a planapo objective (numerical aperture 1.4).

5           An example of transfer of a negatively charged oligoside can be illustrated by the following :

Cells are incubated for 4 h at 37°C with 0.5 mg/ml fluorescein-labelled polyanionic dextrans (either Mw 3000 or Mw 70000) in the absence or in the presence of an oligomeric conjugate. Cells are washed with PBS, fixed with 2 % of p-formaldehyde, washed and mounted on slides in a PBS/glycerol mixture (1:1 v/v) containing 10 mg/ml DABCO (1,4 diazobicyclo-(2,2,2)-octane) as antifading agent. Cells are analyzed with a confocal microscope imaging system (MRC-1024, Bio-Rad) equipped with a Nikon Optiphot epifluorescence microscope (Nikon, Tokyo, Japan) and a planapo objective (numerical aperture 1.4).

15           The fixation of an oligoside to intracellular cofactor can be carried as described in the following example:

An oligoanion with silicated saccharidic components complexed according to the invention is transported into the cytoplasm of human cells where it binds to intracellular cofactors or second messengers such as NF kappa B. This binding causes nuclear transfer of the cofactor which derepress or stimulates genes coding for cytokines (such IL1, TNF- $\alpha$ , IL-12).

This results in a marked stimulation of the human cell cultured in the presence of the complex.

25           The invention also relates to a pharmaceutical composition, comprising as active substance at least an oligomeric conjugate as defined above, or a composition as defined above, or a combined preparation as defined above, or in association with a pharmaceutically acceptable vehicle.

30           The invention also relates to the use of an oligomeric conjugate as defined above, or of a composition as defined above, or of a combined preparation as defined above, or for the preparation of a drug for use in the treatment of cancer, inflammatory or immunology diseases (such as graft rejection, allergy, auto-immunity) or infectious diseases.

The invention also relates to a kit or case containing :

- an oligomeric conjugate as defined above, substituted by a protonable residue leading in a weak acid medium to a destabilization of cellular membranes, this oligomeric conjugate being able to comprise a recognition signal, which is previously fixed or not on the above-said conjugate, said recognition signal being dependent upon the cell to target,

- at least one biological molecule to transfer,  
- optionally reagents enabling the possible binding of the recognition signal on the above-said oligomeric conjugate,

- optionally reagents enabling the formation of a composition as defined above, or of a combined preparation as defined above,

- reagents enabling the transfer of the biological molecule in the cytosol and/or the cell nucleus.

### **LEGEND OF THE FIGURES**

#### *Inhibition of luciferase gene expression by GEM-91.*

**Figure 1** shows the activity of GEM-91, an antisense phosphorothioate oligonucleotide (PS-ODN) (CTC TCG CAC CCA TCT CTC TCC TTC T) complementary to the AUG initiation site of gag HIV-1 gene. The effect of histidylated oligolysines was evaluated by using pRET-Luc cells (a rabbit smooth muscle cell line). These cells produce endogenous luciferase under the control of the human phosphoglycerate kinase promoter and the luciferase gene sequence around the AUG codon was replaced by the initiator AUG codon and several downstream codons of gagHIV-1 gene. The results showed that the activity of GEM-91 ( $IC_{50} > 5 \mu M$ ) was increased more than 10 times in the presence of  $20 \mu M$  HoK2 ( $IC_{50} 0.25 \mu M$ ). Whilst, no significant inhibition was obtained in the presence of HoK3 in which the  $\alpha-NH_2$  histidyl residues were acetylated, suggesting that interactions between ODN and histidylated oligolysines were involved. pRET-Luc cells, seeded onto 24-well plates ( $2 \times 10^5$  cells/well), were treated for 4 h at



37°C in DMEM supplemented with 2 % FBS containing various concentrations of GEM-91, (■) in the absence of histidylated oligolysine, ( ) in the presence of 20 µM HoK2 or (○) in the presence of 20 µM HoK3. HoK2 and HoK3 are histidylated oligolysines prepared as described in the following text. Then, FBS was raised to 6 % and cells were further incubated for 18 h. Luciferase gene expression was measured by recording luminescence for 4 s. The percentage of luciferase inhibition was calculated by using ,  $[(RLU^{ODN}-RLU)/RLU] \times 100$  where  $RLU^{ODN}$  and RLU were the luciferase activity into cell lysates of cells incubated in the absence and in the presence of ODN, respectively. Results shown typical of experiments carried out in triplicate and repeated at least twice. Data are means  $\pm$  standard deviation.

*Inhibition of TNF- $\alpha$  induced ICAM-1 expression by ISI 1939.*

Figure 2 shows the inhibitory effect of TNF- $\alpha$  induced ICAM-1 expression by ISIS 1939 (CCCCCACCACCTTCCCCTCT), an antisense phosphorothioate oligonucleotide (PS-ODN) targeted to the 3' non-coding region of ICAM-1 mRNA. The results showed that TNF- $\alpha$  induced ICAM-1 expression was inhibited by ISIS 1939 in the presence of 20 µM of histidylated oligolysines. HoK2 ( $IC_{50}$  of 0.25 µM) appeared to be more efficient than HoK1 ( $IC_{50}$  of 0.5 µM) probably because HoK2 bore less histidyl residues than HoK1 (15 *versus* 12). The inhibition was very low in the absence of histidylated oligolysines even up to 1 µM ODN (20 % inhibition). A549 cells (ATCC CCL 185, ATCC Rockville, MD) were plated onto 96-wells microtiter plates ( $10^4$  cells /well). The day after, culture medium was removed and cells were washed. Cells were incubated at 37°C for 4 h in 100 µl DMEM serum-free medium containing ISIS 1939 ODN either in the absence (■) or in the presence of 20 µM (●) HoK1 or (□) HoK2. HoK2 and HoK3 are histidylated oligolysines prepared as described in the following text. One volume of fresh medium containing 10 ng/ml TNF- $\alpha$  was added and cells were further incubated for 18 h. ICAM-1 expression was quantified by ELISA using anti-ICAM-1 antibodies. Cells were washed 3 times with 200 µl of PBS and fixed for 20 min at room temperature in PBS containing 20 mg/ml paraformaldehyde. Then, cells were incubated for 90 min at 37°C with anti-ICAM 1 mouse antibody (Becton

Dickinson) diluted 20 times in PBS containing 20 mg/ml BSA. Cells were washed 3 times with PBS and then incubated for 1 h at 37°C with an anti-mouse horseradish peroxidase conjugate (Becton Dickinson) diluted 2000 times in PBS containing 20 mg/ml BSA. After 3 washes, the peroxidase activity was assessed by using 100 µl of o-phenylenediamine dihydrochloride peroxidase substrate tablet set (Sigma). After X min incubation at 37°C, the reaction was stopped by adding 25 µl of 3 N H<sub>2</sub>SO<sub>4</sub> and the absorbance read at 492 nm. All calculations were made relative to untreated controls in the absence or in the presence of TNF-α. The percentage of TNF-α-induced expression of ICAM-1 was calculated as follows :  
5  
10  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65  
66  
67  
68  
69  
70  
71  
72  
73  
74  
75  
76  
77  
78  
79  
80  
81  
82  
83  
84  
85  
86  
87  
88  
89  
90  
91  
92  
93  
94  
95  
96  
97  
98  
99  
100  
101  
102  
103  
104  
105  
106  
107  
108  
109  
110  
111  
112  
113  
114  
115  
116  
117  
118  
119  
120  
121  
122  
123  
124  
125  
126  
127  
128  
129  
130  
131  
132  
133  
134  
135  
136  
137  
138  
139  
140  
141  
142  
143  
144  
145  
146  
147  
148  
149  
150  
151  
152  
153  
154  
155  
156  
157  
158  
159  
160  
161  
162  
163  
164  
165  
166  
167  
168  
169  
170  
171  
172  
173  
174  
175  
176  
177  
178  
179  
180  
181  
182  
183  
184  
185  
186  
187  
188  
189  
190  
191  
192  
193  
194  
195  
196  
197  
198  
199  
200  
201  
202  
203  
204  
205  
206  
207  
208  
209  
210  
211  
212  
213  
214  
215  
216  
217  
218  
219  
220  
221  
222  
223  
224  
225  
226  
227  
228  
229  
230  
231  
232  
233  
234  
235  
236  
237  
238  
239  
240  
241  
242  
243  
244  
245  
246  
247  
248  
249  
250  
251  
252  
253  
254  
255  
256  
257  
258  
259  
260  
261  
262  
263  
264  
265  
266  
267  
268  
269  
270  
271  
272  
273  
274  
275  
276  
277  
278  
279  
280  
281  
282  
283  
284  
285  
286  
287  
288  
289  
290  
291  
292  
293  
294  
295  
296  
297  
298  
299  
300  
301  
302  
303  
304  
305  
306  
307  
308  
309  
310  
311  
312  
313  
314  
315  
316  
317  
318  
319  
320  
321  
322  
323  
324  
325  
326  
327  
328  
329  
330  
331  
332  
333  
334  
335  
336  
337  
338  
339  
340  
341  
342  
343  
344  
345  
346  
347  
348  
349  
350  
351  
352  
353  
354  
355  
356  
357  
358  
359  
360  
361  
362  
363  
364  
365  
366  
367  
368  
369  
370  
371  
372  
373  
374  
375  
376  
377  
378  
379  
380  
381  
382  
383  
384  
385  
386  
387  
388  
389  
390  
391  
392  
393  
394  
395  
396  
397  
398  
399  
400  
401  
402  
403  
404  
405  
406  
407  
408  
409  
410  
411  
412  
413  
414  
415  
416  
417  
418  
419  
420  
421  
422  
423  
424  
425  
426  
427  
428  
429  
430  
431  
432  
433  
434  
435  
436  
437  
438  
439  
440  
441  
442  
443  
444  
445  
446  
447  
448  
449  
450  
451  
452  
453  
454  
455  
456  
457  
458  
459  
460  
461  
462  
463  
464  
465  
466  
467  
468  
469  
470  
471  
472  
473  
474  
475  
476  
477  
478  
479  
480  
481  
482  
483  
484  
485  
486  
487  
488  
489  
490  
491  
492  
493  
494  
495  
496  
497  
498  
499  
500  
501  
502  
503  
504  
505  
506  
507  
508  
509  
510  
511  
512  
513  
514  
515  
516  
517  
518  
519  
520  
521  
522  
523  
524  
525  
526  
527  
528  
529  
530  
531  
532  
533  
534  
535  
536  
537  
538  
539  
540  
541  
542  
543  
544  
545  
546  
547  
548  
549  
550  
551  
552  
553  
554  
555  
556  
557  
558  
559  
560  
561  
562  
563  
564  
565  
566  
567  
568  
569  
570  
571  
572  
573  
574  
575  
576  
577  
578  
579  
580  
581  
582  
583  
584  
585  
586  
587  
588  
589  
590  
591  
592  
593  
594  
595  
596  
597  
598  
599  
600  
601  
602  
603  
604  
605  
606  
607  
608  
609  
610  
611  
612  
613  
614  
615  
616  
617  
618  
619  
620  
621  
622  
623  
624  
625  
626  
627  
628  
629  
630  
631  
632  
633  
634  
635  
636  
637  
638  
639  
640  
641  
642  
643  
644  
645  
646  
647  
648  
649  
650  
651  
652  
653  
654  
655  
656  
657  
658  
659  
660  
661  
662  
663  
664  
665  
666  
667  
668  
669  
670  
671  
672  
673  
674  
675  
676  
677  
678  
679  
680  
681  
682  
683  
684  
685  
686  
687  
688  
689  
690  
691  
692  
693  
694  
695  
696  
697  
698  
699  
700  
701  
702  
703  
704  
705  
706  
707  
708  
709  
710  
711  
712  
713  
714  
715  
716  
717  
718  
719  
720  
721  
722  
723  
724  
725  
726  
727  
728  
729  
730  
731  
732  
733  
734  
735  
736  
737  
738  
739  
740  
741  
742  
743  
744  
745  
746  
747  
748  
749  
750  
751  
752  
753  
754  
755  
756  
757  
758  
759  
760  
761  
762  
763  
764  
765  
766  
767  
768  
769  
770  
771  
772  
773  
774  
775  
776  
777  
778  
779  
780  
781  
782  
783  
784  
785  
786  
787  
788  
789  
790  
791  
792  
793  
794  
795  
796  
797  
798  
799  
800  
801  
802  
803  
804  
805  
806  
807  
808  
809  
810  
811  
812  
813  
814  
815  
816  
817  
818  
819  
820  
821  
822  
823  
824  
825  
826  
827  
828  
829  
830  
831  
832  
833  
834  
835  
836  
837  
838  
839  
840  
841  
842  
843  
844  
845  
846  
847  
848  
849  
850  
851  
852  
853  
854  
855  
856  
857  
858  
859  
860  
861  
862  
863  
864  
865  
866  
867  
868  
869  
870  
871  
872  
873  
874  
875  
876  
877  
878  
879  
880  
881  
882  
883  
884  
885  
886  
887  
888  
889  
890  
891  
892  
893  
894  
895  
896  
897  
898  
899  
900  
901  
902  
903  
904  
905  
906  
907  
908  
909  
910  
911  
912  
913  
914  
915  
916  
917  
918  
919  
920  
921  
922  
923  
924  
925  
926  
927  
928  
929  
930  
931  
932  
933  
934  
935  
936  
937  
938  
939  
940  
941  
942  
943  
944  
945  
946  
947  
948  
949  
950  
951  
952  
953  
954  
955  
956  
957  
958  
959  
960  
961  
962  
963  
964  
965  
966  
967  
968  
969  
970  
971  
972  
973  
974  
975  
976  
977  
978  
979  
980  
981  
982  
983  
984  
985  
986  
987  
988  
989  
990  
991  
992  
993  
994  
995  
996  
997  
998  
999  
1000  
1001  
1002  
1003  
1004  
1005  
1006  
1007  
1008  
1009  
1010  
1011  
1012  
1013  
1014  
1015  
1016  
1017  
1018  
1019  
1020  
1021  
1022  
1023  
1024  
1025  
1026  
1027  
1028  
1029  
1030  
1031  
1032  
1033  
1034  
1035  
1036  
1037  
1038  
1039  
1040  
1041  
1042  
1043  
1044  
1045  
1046  
1047  
1048  
1049  
1050  
1051  
1052  
1053  
1054  
1055  
1056  
1057  
1058  
1059  
1060  
1061  
1062  
1063  
1064  
1065  
1066  
1067  
1068  
1069  
1070  
1071  
1072  
1073  
1074  
1075  
1076  
1077  
1078  
1079  
1080  
1081  
1082  
1083  
1084  
1085  
1086  
1087  
1088  
1089  
1090  
1091  
1092  
1093  
1094  
1095  
1096  
1097  
1098  
1099  
1100  
1101  
1102  
1103  
1104  
1105  
1106  
1107  
1108  
1109  
1110  
1111  
1112  
1113  
1114  
1115  
1116  
1117  
1118  
1119  
1120  
1121  
1122  
1123  
1124  
1125  
1126  
1127  
1128  
1129  
1130  
1131  
1132  
1133  
1134  
1135  
1136  
1137  
1138  
1139  
1140  
1141  
1142  
1143  
1144  
1145  
1146  
1147  
1148  
1149  
1150  
1151  
1152  
1153  
1154  
1155  
1156  
1157  
1158  
1159  
1160  
1161  
1162  
1163  
1164  
1165  
1166  
1167  
1168  
1169  
1170  
1171  
1172  
1173  
1174  
1175  
1176  
1177  
1178  
1179  
1180  
1181  
1182  
1183  
1184  
1185  
1186  
1187  
1188  
1189  
1190  
1191  
1192  
1193  
1194  
1195  
1196  
1197  
1198  
1199  
1200  
1201  
1202  
1203  
1204  
1205  
1206  
1207  
1208  
1209  
1210  
1211  
1212  
1213  
1214  
1215  
1216  
1217  
1218  
1219  
1220  
1221  
1222  
1223  
1224  
1225  
1226  
1227  
1228  
1229  
1230  
1231  
1232  
1233  
1234  
1235  
1236  
1237  
1238  
1239  
1240  
1241  
1242  
1243  
1244  
1245  
1246  
1247  
1248  
1249  
1250  
1251  
1252  
1253  
1254  
1255  
1256  
1257  
1258  
1259  
1260  
1261  
1262  
1263  
1264  
1265  
1266  
1267  
1268  
1269  
1270  
1271  
1272  
1273  
1274  
1275  
1276  
1277  
1278  
1279  
1280  
1281  
1282  
1283  
1284  
1285  
1286  
1287  
1288  
1289  
1290  
1291  
1292  
1293  
1294  
1295  
1296  
1297  
1298  
1299  
1300  
1301  
1302  
1303  
1304  
1305  
1306  
1307  
1308  
1309  
1310  
1311  
1312  
1313  
1314  
1315  
1316  
1317  
1318  
1319  
1320  
1321  
1322  
1323  
1324  
1325  
1326  
1327  
1328  
1329  
1330  
1331  
1332  
1333  
1334  
1335  
1336  
1337  
1338  
1339  
1340  
1341  
1342  
1343  
1344  
1345  
1346  
1347  
1348  
1349  
1350  
1351  
1352  
1353  
1354  
1355  
1356  
1357  
1358  
1359  
1360  
1361  
1362  
1363  
1364  
1365  
1366  
1367  
1368  
1369  
1370  
1371  
1372  
1373  
1374  
1375  
1376  
1377  
1378  
1379  
1380  
1381  
1382  
1383  
1384  
1385  
1386  
1387  
1388  
1389  
1390  
1391  
1392  
1393  
1394  
1395  
1396  
1397  
1398  
1399  
1400  
1401  
1402  
1403  
1404  
1405  
1406  
1407  
1408  
1409  
1410  
1411  
1412  
1413  
1414  
1415  
1416  
1417  
1418  
1419  
1420  
1421  
1422  
1423  
1424  
1425  
1426  
1427  
1428  
1429  
1430  
1431  
1432  
1433  
1434  
1435  
1436  
1437  
1438  
1439  
1440  
1441  
1442  
1443  
1444  
1445  
1446  
1447  
1448  
1449  
1450  
1451  
1452  
1453  
1454  
1455  
1456  
1457  
1458  
1459  
1460  
1461  
1462  
1463  
1464  
1465  
1466  
1467  
1468  
1469  
1470  
1471  
1472  
1473  
1474  
1475  
1476  
1477  
1478  
1479  
1480  
1481  
1482  
1483  
1484  
1485  
1486  
1487  
1488  
1489  
1490  
1491  
1492  
1493  
1494  
1495  
1496  
1497  
1498  
1499  
1500  
1501  
1502  
1503  
1504  
1505  
1506  
1507  
1508  
1509  
1510  
1511  
1512  
1513  
1514  
1515  
1516  
1517  
1518  
1519  
1520  
1521  
1522  
1523  
1524  
1525  
1526  
1527  
1528  
1529  
1530  
1531  
1532  
1533  
1534  
1535  
1536  
1537  
1538  
1539  
1540  
1541  
1542  
1543  
1544  
1545  
1546  
1547  
1548  
1549  
1550  
1551  
1552  
1553  
1554  
1555  
1556  
1557  
1558  
1559  
1560  
1561  
1562  
1563  
1564  
1565  
1566  
1567  
1568  
1569  
1570  
1571  
1572  
1573  
1574  
1575  
1576  
1577  
1578  
1579  
1580  
1581  
1582  
1583  
1584  
1585  
1586  
1587  
1588  
1589  
1590  
1591  
1592  
1593  
1594  
1595  
1596  
1597  
1598  
1599  
1600  
1601  
1602  
1603  
1604  
1605  
1606  
1607  
1608  
1609  
1610  
1611  
1612  
1613  
1614  
1615  
1616  
1617  
1618  
1619  
1620  
1621  
1622  
1623  
1624  
1625  
1626  
1627  
1628  
1629  
1630  
1631  
1632  
1633  
1634  
1635  
1636  
1637  
1638  
1639  
1640  
1641  
1642  
1643  
1644  
1645  
1646  
1647  
1648  
1649  
1650  
1651  
1652  
1653  
1654  
1655  
1656  
1657  
1658  
1659  
1660  
1661  
1662  
1663  
1664  
1665  
1666  
1667  
1668  
1669  
1670  
1671  
1672  
1673  
1674  
1675  
1676  
1677  
1678  
1679  
1680  
1681  
1682  
1683  
1684  
1685  
1686  
1687  
1688  
1689  
1690  
1691  
1692  
1693  
1694  
1695  
1696  
1697  
1698  
1699  
1700  
1701  
1702  
1703  
1704  
1705  
1706  
1707  
1708  
1709  
1710  
1711  
1712  
1713  
1714  
1715  
1716  
1717  
1718  
1719  
1720  
1721  
1722  
1723  
1724  
1725  
1726  
1727  
1728  
1729  
1730  
1731  
1732  
1733  
1734  
1735  
1736  
1737  
1738  
1739  
1740  
1741  
1742  
1743  
1744  
1745  
1746  
1747  
1748  
1749  
1750  
1751  
1752  
1753  
1754  
1755  
1756  
1757  
1758  
1759  
1760  
1761  
1762  
1763  
1764  
1765  
1766  
1767  
1768  
1769  
1770  
1771  
1772  
1773  
1774  
1775  
1776  
1777  
1778  
1779  
1780  
1781  
1782  
1783  
1784  
1785  
1786  
1787  
1788  
1789  
1790  
1791  
1792  
1793  
1794  
1795  
1796  
1797  
1798  
1799  
1800  
1801  
1802  
1803  
1804  
1805  
1806  
1807  
1808  
1809  
1810  
1811  
1812  
1813  
1814  
1815  
1816  
1817  
1818  
1819  
1820  
1821  
1822  
1823  
1824  
1825  
1826  
1827  
1828  
1829  
1830  
1831  
1832  
1833  
1834  
1835  
1836  
1837  
1838  
1839  
1840  
1841  
1842  
1843  
1844  
1845  
1846  
1847  
1848  
1849  
1850  
1851  
1852  
1853  
1854  
1855  
1856  
1857  
1858  
1859  
1860  
1861  
1862  
1863  
1864  
1865  
1866  
1867  
1868  
1869  
1870  
1871  
1872  
1873  
1874  
1875  
1876  
1877  
1878  
1879  
1880  
1881  
1882  
1883  
1884  
1885  
1886  
1887  
1888  
1889  
1890  
1891  
1892  
1893  
1894  
1895  
1896  
1897  
1898  
1899  
1900  
1901  
1902  
1903  
1904  
1905  
1906  
1907  
1908  
1909  
1910  
1911  
1912  
1913  
1914  
1915  
1916  
1917  
1918  
1919  
1920  
1921  
1922  
1923  
1924  
1925  
1926  
1927  
1928  
1929  
1930  
1931  
1932  
1933  
1934  
1935  
1936  
1937  
1938  
1939  
1940  
1941  
1942  
1943  
1944  
1945  
1946  
1947  
1948  
1949  
1950  
1951  
1952  
1953  
1954  
1955  
1956  
1957  
1958  
1959  
1960  
1961  
1962  
1963  
1964  
1965  
1966  
1967  
1968  
1969  
1970  
1971  
1972  
1973  
1974  
1975  
1976  
1977  
1978  
1979  
1980  
1981  
1982  
1983  
1984  
1985  
1986  
1987  
1988  
1989  
1990  
1991  
1992  
1993  
1994  
1995  
1996  
1997  
1998  
1999  
2000  
2001  
2002  
2003  
2004  
2005  
2006  
2007  
2008  
2009  
2010  
2011  
2012  
2013  
2014  
2015  
2016  
2017  
2018  
2019  
2020  
2021  
2022  
2023  
2024  
2025  
2026  
2027  
2028  
2029  
2030  
2031  
2032  
2033  
2034  
2035  
2036  
2037  
2038  
2039  
2040  
2041  
2042  
2043  
2044  
2045  
2046  
2047  
2048  
2049  
2050  
2051  
2052  
2053  
2054  
2055  
2056  
2057  
2058  
2059  
2060  
2061  
2062  
2063  
2064  
2065  
2066  
2067  
2068  
2069  
2070  
2071  
2072  
2073  
2074  
2075  
2076  
2077  
2078  
2079  
2080  
2081  
2082  
2083  
2084  
2085  
2086  
2087  
2088  
2089  
2090  
2091  
2092  
2093  
2094  
2095  
2096  
2097  
2098  
2099  
2100  
2101  
2102  
2103  
2104  
2105  
2106  
2107  
2108  
2109  
2110  
2111  
2112  
2113  
2114  
2115  
2116  
2117  
2118  
2119  
2120  
2121  
2122  
2123  
2124  
2125  
2126  
2127  
2128  
2129  
2130  
2131  
2132  
2133  
2134  
2135  
2136  
2137  
2138  
2139  
2140  
2141  
2142  
2143  
2144  
2145  
2146  
2147  
2148  
2149  
2150  
2151  
2152  
2153  
2154  
2155  
2156  
2157  
2158  
2159  
2160  
2161  
2162  
2163  
2164  
2165  
2166  
2167  
2168  
2169  
2170  
2171  
2172  
2173  
2174  
2175  
2176  
2177  
2178  
2179  
2180  
2181  
2182  
2183  
2184  
2185  
2186  
2187  
2188  
2189  
2190  
2191  
2192  
2193  
2194  
2195  
2196

ATCC Rockville, MD) were seeded onto sterile coverslips in 20-mm wells ( $2 \times 10^5$  cells/ well) and allowed to adhere. Cells were incubated in the presence of 0.125  $\mu$ M fluorescein-labelled PS-ODN for 4 h at 37°C (a) in the absence or in the presence of 20  $\mu$ M (b) HoK1, (c) HoK2, (d) HoK3 or (e) Plk (Oligolysine containing 19 lysyl residues and non substituted by histidine used as control). Cells were fixed with 2 % of p-formaldehyde, washed and mounted on slides in a PBS/glycerol mixture (1:1 v/v) containing 10 mg/ml DABCO (1,4 diazobicyclo-(2,2,2)-octane) as antifading agent. Cells were analyzed with a confocal microscope imaging system (MRC-1024, Bio-Rad) equipped with a Nikon Optiphot epifluorescence microscope (Nikon, Tokyo, Japan) and a planapo objective (numerical aperture 1.4).

## EXAMPLES

### *Preparation of histidylated oligolysines :*

Oligolysine (Poly-L-lysine, HBr ; average molecular weight of 3950 ; average degree of polymerization of 19) (Bachem Feinchemikalien, Bubendorf, Switzerland) (1 g in 200 ml H<sub>2</sub>O) was passed through an anion exchange column (Dowex 2 x 8, OH form, 20-50 mesh) in order to remove bromide ions. The eluate was neutralized with a 10 % *p*-toluene sulfonic acid solution in water and freeze-dried.

### Example 1 : Preparation of HoK1

Oligolysine *p*-toluene sulfonate salt (50 mg ; 8.6  $\mu$ mol) in 2 ml dimethylsulfoxide (Aldrich, Strasbourg, France) in the presence of diisopropylethylamine (50  $\mu$ l ; 344  $\mu$ mol) (Aldrich) was reacted for 20 h at 20°C with (Boc)His(Boc)-OH (64 mg ; 146  $\mu$ mol) (Novabiochem, Bad Soden, Germany) in the presence of benzotriazol-1-yl-oxy-tris-(dimethylamino) phosphonium hexafluorophosphate (BOP) (Richelieu Biotechnologies, Saint Hyacinthe, Canada) (159mg ; 358  $\mu$ mol). The residual  $\epsilon$ -amino groups of oligolysine was then substituted with gluconoyl residues (GlcA) :  $\delta$ -gluconolactone (86mg ; 48  $\mu$ mol)

(Aldrich) and diisopropylethylamine (134  $\mu$ l ; 921  $\mu$ mol) were added and the solution was stirred for 20h at 20°C. The *N*-protecting Boc groups were removed by acidic treatment by adding 10 volumes of a H<sub>2</sub>O/trifluoroacetic acid mixture (1:1 ; v/v) for 24h at 20°C. Water and trifluoroacetic acid were removed under reduced pressure. HoK1 was precipitated by adding 10 volumes of isopropanol and spun down by centrifugation (1800g for 15 min). The pellet was washed with isopropanol, collected by centrifugation (1800g for 15 min), solubilized in distilled water and freeze-dried. The average number of histidyl residues bound per oligolysine molecule was determined by <sup>1</sup>H-NMR spectroscopy at 300 MHz in D<sub>2</sub>O according to :  $x = 6 \cdot (h_{8.7} / h_{Lys}) \cdot DP$ , where  $h_{8.7}$  was the value of the integration of the signal at 8.7 ppm corresponding to the proton (1H C<sub>12</sub>) of histidyl residues,  $h_{Lys}$  that in the range from 1.3 to 1.9 ppm corresponding to the 6 methylene protons (C<sub>3</sub>, C<sub>4</sub> and C<sub>5</sub>) of lysyl residues and DP the degree of polymerization of oligolysine. The number of histidyl residues bound per oligolysine molecule was 12. The average number of gluconoyl residues bound per oligolysine molecule was determined by <sup>1</sup>H-NMR spectroscopy from :  $x = 3/2 \cdot (h_G / h_{Lys}) \cdot DP$ , where  $h_G$  was the value of the integration in the range 3.6 to 3.9 ppm of the 4 protons (1H C<sub>10</sub>, 1H C<sub>11</sub> and 2H C<sub>12</sub>) of gluconoyl residues,  $h_{Lys}$  that in the range of 1.3 to 1.9 ppm of the 6 methylene protons (C<sub>3</sub>, C<sub>4</sub> and C<sub>5</sub>) of lysyl residues and DP the degree of polymerization of pLK. The number of gluconoyl residues bound per oligolysine molecule was 3. The number of free  $\epsilon$ -amino groups per oligolysine molecule was 4.

#### Example 2 : Preparation of HoK2

Oligolysine *p*-toluene sulfonate salt (80 mg ; 13.7  $\mu$ mol) in 3 ml dimethylsulfoxide in the presence of diisopropylethylamine (90  $\mu$ l ; 620  $\mu$ mol) was reacted for 6 h at 20°C with the *N*-hydroxysuccinimidyl derivative of (Boc)His(Boc)-OH (92 mg ; 204  $\mu$ mol) (Bachem Feinchemikalien). The residual  $\epsilon$ -amino groups of oligolysine was then acetylated (Ac) : acetic anhydride (13  $\mu$ l ; 109  $\mu$ mol) (Aldrich) and diisopropylethylamine (5  $\mu$ l ; 35  $\mu$ mol) were added and the solution was stirred for 30 min at 20°C. The *N*-protecting Boc groups were removed by acidic treatment by adding 10 volumes of a


H<sub>2</sub>O/trifluoroacetic acid mixture (1 : 1 ; v/v) for 2 h at 20°C. Water and trifluoroacetic acid were removed under reduced pressure. HoK2 was precipitated by adding 10 volumes of isopropanol and spun down by centrifugation (1800 g for 15 min). The pellet was washed with isopropanol, collected by centrifugation (1800 g for 15 min), solubilized in distilled water and freeze-dried. The average number of histidyl residues bound per oligolysine molecule was determined by <sup>1</sup>H-NMR spectroscopy at 300 MHz in D<sub>2</sub>O according to :  $x = 6 \cdot (h_{8.7} / h_{lys})$ . DP, where  $h_{8.7}$  was the value of the integration of the signal at 8.7 ppm corresponding to the proton (1H C<sub>12</sub>) of histidyl residues,  $h_{lys}$  that in the range from 1.3 to 1.9 ppm corresponding to the 6 methylene protons (C<sub>3</sub>, C<sub>4</sub> and C<sub>5</sub>) of lysyl residues and DP the degree of polymerization of oligolysine. The number of histidyl residues bound per oligolysine molecule was 15. The average number of acetyl residues bound per oligolysine molecule was determined by <sup>1</sup>H-NMR spectroscopy from :  $x = (h_A / h_{lys}) \cdot DP$ , where  $h_A$  was the value of the integration at 2.04 ppm of the 3 protons of acetyl residues,  $h_{lys}$  that in the range of 1.3 to 1.9 ppm of the 6 methylene protons (C<sub>3</sub>, C<sub>4</sub> and C<sub>5</sub>) of lysyl residues and DP the degree of polymerization of pLK. The number of acetyl residues bound per oligolysine molecule was 3. The number of free ε-amino groups per oligolysinemolecule was 1.

### Example 3 : Preparation of HoK3

Oligolysine *p*-toluene sulfonate salt (85 mg ; 14.6 μmol) in 3 ml dimethylsulfoxide in the presence of diisopropylethylamine (80 μl ; 555 μmol) was reacted for 20 h at 20°C with N-acetyl-His-OH (288 mg ; 307 μmol) (Sigma) in the presence of BOP (265 mg ; 597 μmol). The residual ε-amino groups of oligolysine was then acetylated (Ac) with acetic anhydride for 30 min at 20°C. HoK3 was precipitated by adding 10 volumes of isopropanol and spun down by centrifugation (1800 g for 15 min). The pellet was washed with isopropanol, collected by centrifugation (1800 g for 15 min), solubilized in distilled water and freeze-dried. The average number of histidyl residues bound per oligolysine molecule was determined by <sup>1</sup>H-NMR spectroscopy as describe The number of

histidyl residues bound per oligolysine molecule was 15. The number of free  $\epsilon$ -amino groups per oligolysine molecule was 1.

Example 4 : Preparation of synthons for synthesis of oligomeric conjugates



wherein

R and R' represent aminoprotecting groups,  $a_i$  is an integer varying from 0 to 6,  $b_i$  is an integer varying from 0 to 6,  $n$  is an integer varying from 1 to 6,  $n'$  is an integer varying from 0 to 6,  $n''$  is an integer varying from 0 to 6,  $m$  is an integer varying from 1 to 6.

An example of synthon : the Lys(His) synthon

wherein

$$a_i = b_i = 0$$

$$n = 4$$

$$R_1 = \text{Fmoc}$$

$$n' = n'' = 0$$

$$m = 1$$

$$R' = \text{Boc}$$

$$B = (\text{NBoc})\text{Imidazole}$$

The N-hydroxysuccinimidyl derivative of (Boc)<sub>2</sub>-His-OH [(Boc)<sub>2</sub>-His-OSu] (1g ; 2.2 mmol) is coupled to Fmoc-Lys-OH (722 mg ; 2.2 mmol) for 24 h at 20°C in dimethylformamide (ml). The Lys(His) synthon is precipitated with isopropanol, collected by centrifugation, washed with ether and dried under vacuum. The Lys(His) synthon is purified by cristallization.

Example 5 :

Preparation of an histidylated oligolysine (HoK) containing 20 lysine residues and 20 histidyl residues.

A HoK containing exactly 20 lysyl residues and 20 histidyl residues can be entirely synthetised by using the above Lys (His) synthon. Briefly, 20 Lys(His) synthon are successively assembled on a Applied Biosystems 433A synthesizer with conductimetric monitoring by using Fmoc-protected amino acids. Lys(His) synthons are coupled by the HBTU activation method. HoK are cleaved from the resin and side chain protecting Boc groups are removed with a trifluoroacetic acid/water mixture (50% : 50% ; v/v) for 3 h at room temperature. Crude HoK is precipitated with isopropanol and collected by centrifugation. HoK is washed three times with isopropanol, resuspended in distilled water and freeze-dried.

Example 6

Preparation of an oligomeric conjugate containing 17 lysyl residues substituted with 17 histidyl residues and 3 leucyl residues inserted anywhere in the lys (His) sequence

Oligomers made of exactly 17 lysyl residues substituted with 17 histidyl residues and 3 leucyl residues inserted anywhere in the Lys(His) sequence can be entirely synthetised by using the above Lys (His) synthon and Fmoc Leu on a Applied Biosystems 433A synthesizer with conductimetric monitoring by using Fmoc-protected amino acids. Lys(His) synthons and Leu were coupled by the HBTU activation method. Oligomers are cleaved from the resin and side chain protecting Boc groups are removed with a trifluoroacetic acid/water mixture (50% : 50% ; v/v) for 3 h at room temperature. Oligomers are precipitated with isopropanol and collected by centrifugation. Oligomers are washed three times with isopropanol, resuspended in distilled water and freeze-dried.

**Example 7 : Preparation of (K(His)-KL(His)-L)<sub>7</sub>**

An oligomer (K(His)-K(His)-L)<sub>7</sub> can be entirely synthesised by using the above Lys (His) synthon and Fmoc Leu on a Applied Biosystems 433A synthesizer with conductimetric monitoring by using Fmoc-protected amino acids. Lys(His) synthons and Leu are coupled by the HBTU activation method. The oligomer (K(His)-K(His)-L)<sub>7</sub> is cleaved from the resin and side chain protecting Boc groups are removed with a trifluoroacetic acid/water mixture (50% : 50% ; v/v) for 3 h at room temperature. The polymer is precipitated with isopropanol and collected by centrifugation. The oligomer is washed three times with isopropanol, resuspended in distilled water and freeze-dried.

**Example 8 : Preparation of (K(His)-L-K(His))<sub>7</sub>**

A oligomer (K(His)-L-K(His))<sub>7</sub> can be entirely synthesised by using the above Lys (His) synthon and Fmoc Leu on a Applied Biosystems 433A synthesizer with conductimetric monitoring by using Fmoc-protected amino acids. Lys(His) synthons and Leu are coupled by the HBTU activation method. The oligomer is cleaved from the resin and side chain protecting Boc groups are removed with a trifluoroacetic acid/water mixture (50% : 50% ; v/v) for 3 h at room temperature. The oligomer is precipitated with isopropanol and collected by centrifugation. The oligomer is washed three times with isopropanol, resuspended in distilled water and freeze-dried.

Table I shows that oligolysine having a DP of either 190, 72 or 36 substituted with less than 45% of histidyl residues, do not allow the transfer of small nucleic acid, particularly oligonucleotides. Indeed, it is necessary to adapt the size of histidylated oligolysines. Conversely, small histidylated oligolysines (DP of 36 or 19) substituted by more than 50% of histidyl residues do not allow efficient gene transfer by histidylated oligolysine/plasmid complexes (Table I). In addition oligolysines substituted with less than 50% histidyl residues are cytotoxic.



**Table I** : comparative evaluation of gene transfer and oligonucleotide transfer by using histidylated polylysine.

DP	His (%)	DNA	ODN	Cytotoxicity (%)
190	35	100	0	24
190	45	110	0	21
72	23	87-96	0	24
36	22	61-100	0	25
36	53	0-10	20	4
19	25	15-26	0	49
19	45	9	0	40
19	60	nd	100	26
19	80	nd	100	0
19	100	nd	100	0

DP is the oligolysine degree of polymerization. DNA corresponds to transfection by using histidylated oligolysine/pCMVLUC. The transfection efficiency is scored on a 0 to 100 scale. The transfection efficiency is determined from the luciferase activity in cells measured by luminescence. ODN corresponds to cytosolic and nuclear transfer of fluorescein-labelled oligonucleotide in the presence of histidylated oligolysine, evaluated under confocal microscope. Cytotoxicity was evaluated by using the colorimetric MTT assay. (MTT = 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide).

## CLAIMS

1. Oligomeric conjugate positively charged, containing an oligomer with a  
5 polymerization degree (PD) from 5 to 50, preferably 10 to 40 and more preferably  
20, formed from monomeric components having free  $\text{NH}_3^+$  in a number equal to  
or higher than 50 % of the polymerization degree,

said oligomer being as follows :

10 - the free  $\text{NH}_3^+$  of the above-mentioned components are substituted in a  
ratio of at least 50 %, advantageously from 60 % to 95 %, particularly 80 to 90 %  
(this ratio being determined by nuclear magnetic resonance), by protonable  
residues in a weak acid medium, leading in such a weak acid medium to a  
destabilization of cellular membranes,

15 - the above-mentioned protonable residues possess in addition the following  
properties :

→ they contain a functional group enabling them to be linked to the  
above-mentioned oligomer,

→ they do not correspond to a recognition signal recognized by a  
cellular membrane receptor,

20 → they can comprise at least one free  $\text{NH}_3^+$  group,

- the free  $\text{NH}_3^+$  of the above-mentioned monomers can be also substituted  
by an uncharged residues leading to a reduction of the number of positive charges  
in comparison to the same oligomeric before substitution,

25 - molecules constituting a recognition signal recognized by a membrane  
cellular receptor may be present :

→ either by substitution of some of the free  $\text{NH}_3^+$  of the above-  
mentioned monomers,

→ either on some of the uncharged residues leading to a reduction  
of the number of charges,

30 → either on some of the above-mentioned protonable residues  
leading to a destabilization of the cellular membranes,

→ or by substitution of the free  $\text{NH}_3^+$  (if it is present) of the above-mentioned protonable residues leading to a destabilization of the cellular membrane,

provided that :

1) the total number of the non substituted  $\text{NH}_3^+$  functions is of at least 50 % of the polymerization degree,

2) the number of monomers initially carrying free  $\text{NH}_3^+$  is substituted in a ratio of at least 50 % of the polymerization degree by residues leading to a destabilization of the cellular membrane.

2. Oligomeric conjugate according to claim 1, wherein the protonable residues leading to a destabilization of cellular membranes, present the additional properties :

- they are weak bases the pK of which in aqueous medium is lower than 8, so that a proportion higher than 50 % of these bases linked to a cationic oligomer is not protonated in a neutral medium of pH 7.4.

3. Oligomeric conjugate complex according to claim 1, wherein the protonable residues leading to a destabilization of cellular membranes, present the additional properties:

- they belong to the group of compounds comprising an imidazole ring,
- they belong to the group of quinolins,
- they belong to the group of pterins,
- they belong to the groupe of pyridins.


4. Oligomeric conjugate according to anyone of claims 1 to 3, wherein the protonable residues leading to a destabilization of the cellular membranes are :

- alkylimidazoles in which the alkyl radical comprises from 1 to 10, particularly from 2 to 6 carbon atoms, and in which only one of the nitrogen atoms of the imidazole ring is substituted.

5. Oligomeric conjugate according to anyone of claims 1 to 4, wherein the protonable residues leading to a destabilization of cellular membranes are chosen from :

5 histidine, 4-carboxymethyl-imidazole,  
 3-(1-methyl-imidazol-4yl)-alanine, 3-(3-methyl-imidazol-4yl)-alanine,  
 2-carboxy-imidazole, histamine, 3-imidazol-4yl)-L-lactic acid,  
 2-(1-methyl-imidazol-4yl)ethylamine, 2-(3-methyl-imidazol-4yl)ethylamine,  
 β-alanyl-histidine-(carnosine), 7-chloro-4(amino-1-methylbutylamino)-quinoline,  
 10 N<sup>4</sup>-(7-chloro-4-quinolinyl)-1,4-pentanediamine,  
 8-(4-amino-1-methylbutylamino)-6-methoxy-quinoline (primaquine),  
 N<sup>4</sup>-(6-methoxy-8-quinolinyl)-1,4-pentanediamine, quininic acid,  
 quinoline carboxylic acid, pteric acid, nicotinic acid, quinolinic acid.

15 6. Oligomeric conjugate according to anyone of claims 1 to 5, wherein the oligomeric conjugate contains an oligomer of the following formula :



wherein \* ai is an integer varying from 0 to 10,



\* bi is an integer varying from 0 to 10,

\* i = degree of polymerization from 5 to 50, and

particularly 10 to 40, and preferably 20,



\* n = is an integer varying from 1 to 6, and preferably 4,

\* R represents in a ratio of 50 % to 100 % (corresponding to a number u)





B

or



B

or



B

m is an integer varying from 1 to 6,


n' is an integer varying from 0 to 6,

n'' is an integer varying from 0 to 6,

B is a weak base as defined according to anyone of claims 2 to 4,


R' represents  $\text{NH}_3^+$  (corresponding to a number p),

or NH (corresponding to a number q) substituted by



r being an integer from 1 to 15,



and preferably 1 to 7.



being an integer from 1 to 15, and

preferably 1 to 7, and s being an

integer from 1 to 6, and preferably 6



Flu being a fluorescent molecule

\* R represents in a ratio of 0 % to 50 % (corresponding to  $f$ :  $0 < f \leq u$ )

- $\text{NH}_3^+$  (corresponding to a number  $j$ ),
- NH (corresponding to a number  $k$ ), substituted by


5 -CO-CH<sub>3</sub>

-CO-(CHOH)<sub>r</sub>H

$r$  being an integer from 1 to 15,  
and preferably 1 to 7

10 -CO-(CH<sub>2</sub>)<sub>s</sub>-(CHOH)<sub>r</sub>H

$r$  being an integer from 1 to 15, and  
preferably 1 to 7, and  $s$  being an  
integer from 1 to 6, and preferably 6



-SO<sub>2</sub>-Flu

15 -CO-Flu

-CS-NH-Flu


Flu being a fluorescent molecule

- H (corresponding to a number  $h$ )
- (CH<sub>2</sub>)<sub>n</sub>H,  $n$  being an integer from 1 to 6

(corresponding to a number  $h$ )

- (CH<sub>2</sub>)<sub>n</sub>-OH  $n$  being an integer from 1 to 6

(corresponding to a number  $h$ )

- (CH<sub>2</sub>)<sub>n</sub>-SA' A' = H, CH<sub>3</sub> or s-

(corresponding to a number  $h$ )  $n$  being integer from 1 to 6

with .  $i = u + j + k + h$

. total number of  $\alpha \text{NH}_3^+$  =  $p = u - q$

. total number of  $\omega \text{NH}_3^+$  =  $j = f - (k + h)$

. total number of  $\text{NH}_3^+$  =  $m = p + j + 1$

with the proviso that :

$$1) u \geq i/2$$


$$2) m \geq i/2$$

5

7. Oligomeric conjugate according to anyone of claims 1 to 6, wherein the oligomeric conjugate contains an oligomer of the following formula :

10

15



20

wherein


\* i = degree of polymerization from 5 to 50, and particularly 10 to 40, and preferably 20,

\* n = is an integer varying from 1 to 6, and preferably 4,

25


\* R represents in a ratio of 50 % to 100 % (corresponding to u)

30




or

35



or



m is an integer varying from 1 to 6,

n' is an integer varying from 0 to 6,

n'' is an integer varying from 0 to 6,

B is a weak base as defined according to anyone of claims 2 to 4,

R' represents  $\text{NH}_3^+$  (corresponding to a number p),

or NH (corresponding to a number q) substituted by

-CO-CH<sub>3</sub>

-CO-(CHOH)rH

r being an integer from 1 to 15,


and preferably 1 to 7

-CO-(CH<sub>2</sub>)s-(CHOH)rH

r being an integer from 1 to 15, and

preferably 1 to 7, and s being an

integer from 1 to 6, and preferably 6



-SO<sub>2</sub>-Flu

-CO-Flu

-CS-NH-Flu

Flu being a fluorescent molecule

\* R represents in a ratio of 0 % to 50 % (corresponding to f : 0 < f ≤ 1)

-  $\text{NH}_3^+$  (corresponding to a number j),

- NH (corresponding to a number k), substituted by


-CO-CH<sub>3</sub>

-CO-(CHOH)rH

r being an integer from 1 to 15,


and preferably 1 to 7





r being an integer from 1 to 15, and preferably 1 to 7, and s being an integer from 1 to 6, and preferably 6

5



10

Flu being a fluorescent molecule

- H (corresponding to a number h)

-  $(\text{CH}_2)_n\text{H}$ ,

n being an integer from 1 to 6

(corresponding to a number h)

15


-  $(\text{CH}_2)_n\text{OH}$

n being an integer from 1 to 6

(corresponding to a number h)

-  $(\text{CH}_2)_n\text{SA}'$

A' = H, CH<sub>3</sub> or



(corresponding to a number h)

n being integer from 1 to 6

20

with .  $i = u + j + k + h$

. total number of  $\alpha \text{NH}_3^+ = p = u - q$

. total number of  $\omega \text{NH}_3^+ = j = f - (k + h)$

. total number of  $\text{NH}_3^+ = m = p + j + 1$

25

with the proviso that :


1)  $u \geq i/2$

2)  $m \geq i/2$

30

8. Oligomeric conjugate according to anyone of claims 1 to 7, wherein the oligomeric conjugate contains an oligomer of the formula according to claim 7, wherein

$$i = 19 \quad n = 4$$




wherein

$$n' = n'' = 0$$

$$\text{R}' = \text{NH}_3^+$$

$$m = 1$$

$$\text{B} = \text{imidazole}$$



$$(f) \text{ R} = \text{NH}_3^+$$


$$u = 12$$

$$j = 7$$

5

or

$$i = 19 \quad n = 4$$





wherein

$$n' = n'' = 0$$

$$\text{R}' = \text{NH}_3^+$$

$$m = 1$$

$$\text{B} = \text{imidazole}$$




$$(f) \text{ R} = \text{NH}_3^+$$

$$u = 16$$

$$j = 3$$

or

$$i = 19 \quad n = 4$$






wherein

$$n' = n'' = 0$$

$$\text{R}' = \text{NH}_3^+$$

$$m = 1$$

$$\text{B} = \text{imidazole}$$




$$u = 19$$

$$j = 0$$

5

or

$$i = 19 \quad n = 4$$





wherein

$$n' = n'' = 0$$

$$\text{R}' = \text{NH}_3^+$$

$$m = 1$$

$$\text{B} = \text{imidazole}$$




$$u = 11$$

$$k = 8$$

or

$$i = 19$$

$$n = 4$$




wherein

$$n' = n'' = 0$$

$$\text{R}' = \text{NH}_3^+$$


$$m = 1$$

$$\text{B} = \text{imidazole}$$



$$u = 15$$

$$k = 4$$




5

or

$$i = 19$$

$$n = 4$$




wherein

$$n' = n'' = 0$$

$$\text{R}' = \text{NH}_3^+$$

$$m = 1$$


$$\text{B} = \text{imidazole}$$



$$r = 5$$


$$u = 12$$

$$k = 3$$



or

$$i = 19 \quad n = 4$$



wherein

$$(q) \quad n' = n'' = 20$$

$$\text{R}' = \text{NH-CO-CH}_3$$

$$m = 1$$

$$\text{B} = \text{imidazole}$$


$$\text{R} = \text{NH-CO-CH-NH-CO-CH}_3$$

$$(f) \text{ R} = \text{NH}^+_3$$

$$u = 16$$

$$f = 4$$

$$k = 3$$



5            9. Composition containing at least one oligomeric conjugate according to anyone of claims 1 to 8, in association with at least one biological molecule, such as a peptide, an oligoside or an oligonucleotide or a mixture thereof.

10           10. Combined preparation containing as active substance the following individual components, in the form of a kit-of-parts :

- an oligomeric conjugate according to anyone of claims 1 to 8,
- at least one oligomeric conjugate according to anyone of claims 1 to 8, in association with at least one biological molecule, such as a peptide, an oligoside or an oligonucleotide, or a mixture thereof,

15           for the simultaneous, separate or sequential use, for the *in vitro*, the *in vivo* or the *ex vivo* transfer of said biological molecules into the cytosol and/or cell nucleus.

11. Use of an oligomeric conjugate according to anyone of claims 1 to 8, for the *in vitro*, the *ex vivo* or the *in vivo* intracellular transfer of biological molecules into the cytosol and/or in the cell nucleus.

12. Use of an oligomeric conjugate according to anyone of claims 1 to 8 or of a composition according to claim 9, or of a combined preparation according to claim 10, for the intracellular the *in vitro*, the *ex vivo* or the *in vivo* transfer of a peptide, an oligoside or an oligonucleotide, or a mixture thereof, into the cytosol or/and in the cell nucleus.

13. Use of an oligomeric conjugate according to anyone of claims 1 to 8 or of a composition according to claim 9, or of a combined preparation according to claim 10, wherein the cells are chosen among muscular, epithelial, endothelial, myeloid cells such as monocytes, macrophages and fibroblasts, leukocytes and granulocytes, osteoblasts as well as dendritic cells, stem cells, neuronal cells, or dermal cells.

14. Method for the *in vivo*, the *in vitro* or the *ex vivo* transfer of an oligonucleotide, wherein an oligonucleotide and an oligomeric conjugate according to anyone of claims 1 to 8 or of a composition according to claim 9, or of a combined preparation according to claim 10, are(is) contacted with a medium containing cells to be transferred, under conditions such that there is :

- transfer of an antisense oligonucleotide in the cytosol and/or the cell nucleus where it binds and blocks the complementary mRNA sequence,
- or transfer of an oligonucleotide as activator into the cytosol where it depresses or activates a second messenger in the cytosol, or the corresponding gene in the nucleus,
- or transfer into the cytosol and/or the cell nucleus of oligonucleotides corresponding to a repetitive bacterial type DNA sequence with stimulating or immunodepressive activity,

- or transfer of an oligonucleotide in the cell nucleus where it binds to DNA and forms a triple helix leading to the inhibition of gene expression.

- or transfer into the cytosol and/or the cell nucleus of RNA oligonucleotide acting as decoys which inhibit gene expression by blocking the binding of regulatory factors to the authentic DNA region.

- or transfer into the cytosol and/or the cell nucleus of ribozymes (RNA oligonucleotides) which inhibit gene expression by cleaving the mRNA.

15. Method for the *in vivo*, the *in vitro* or the *ex vivo* transfer of peptide, wherein a peptide and an oligomeric conjugate according to anyone of claims 1 to 8 or of a composition according to claim 9, or of a combined preparation according to claim 10, in particular wherein the peptide is an antigenic peptide, are(is) contacted with a medium containing cells to be transferred, under conditions such that there is a transfer of said antigenic peptide in the cytosol of antigen presenting cells (macrophages, dendritic cells and B cells) where they are processed in proteosomes in order to bind to MHCI molecules, allowing the presentation of the antigenic epitope fixed on MHCI.

16. Method for the *in vivo*, the *in vitro* or the *ex vivo* transfer of an oligoside, wherein an oligoside and an oligomeric conjugate according to anyone of claims 1 to 8 or of a composition according to claim 9, or of a combined preparation according to claim 10, are(is) contacted with a medium containing cells to be transferred, under conditions such that there is a transfer of said oligoside into the cytosol and/or the cell nucleus.

17. Pharmaceutical composition, comprising as active substance at least an oligomeric conjugate according to anyone of claims 1 to 8, or of a composition according to claim 9, or of a combined preparation according to claim 10, or in association with a pharmaceutically acceptable vehicle.

18. Use of an oligomeric conjugate according to anyone of claims 1 to 8, or of a composition according to claim 9, or of a combined preparation according to claim 10, or for the preparation of a drug for use in the treatment of cancer, inflammatory or immunology diseases (such as graft rejection, allergy, auto-immunity) or infectious diseases.

19. Kit or case containing :

- an oligomeric conjugate according to anyone of claims 1 to 8, substituted by a protonable residue leading in a weak acid medium to a destabilization of cellular membranes, this oligomeric conjugate being able to comprise a recognition signal, which is previously fixed or not on the above-said conjugate, said recognition signal being dependent upon the cell to target,

- at least one biological molecule to transfer,

- optionally reagents enabling the possible binding of the recognition signal on the above-said oligomeric conjugate,

- optionally reagents enabling the formation of a composition according to claim 9, or of a combined preparation according to claim 10,

- reagents enabling the transfer of the biological molecule in the cytosol and/or the cell nucleus.



1 / 3




Figure 1

2 / 3




Figure 2

3 / 3




Figure 3

# INTERNATIONAL SEARCH REPORT

International Application No.

PCT/EP 99/08980

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C12N15/11 A61K47/48 A61K48/00

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 98 22610 A (IDM IMMUNO DESIGNED MOLECULES ;MIDOUX PATRICK (FR); MONSIGNY MICHE) 28 May 1998 (1998-05-28) cited in the application * see in particular the claims; and Figures 1,5 *	1-7,9-19
A	WO 98 19710 A (SCHACHT ETIENNE HONORE ;ULBRICH KAREL (CZ); SEYMOUR LEONARD CHARLE) 14 May 1998 (1998-05-14) *see p.39,1.30 - p.40; claims 1,6-8*	1-19
A	US 5 627 270 A (HATZENBUHLER NICOLE T ET AL) 6 May 1997 (1997-05-06) * see claims 1,3; col.71,1.36-38 *	1-19
	-/-	

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

### \* Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "S" document member of the same patent family

Date of the actual completion of the international search

2 March 2000

Date of mailing of the international search report

117.03.00

Name and mailing address of the ISA

European Patent Office, P.B. 6818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3018

Authorized officer

Isert, B

# INTERNATIONAL SEARCH REPORT

Inter. Appl. No.  
PCT/EP 99/08980

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	EP 0 727 223 A (HISAMITSU PHARMACEUTICAL CO) 21 August 1996 (1996-08-21) * see the claims; Figures 8-10; page 7 *	1-19
A	GOTO T. ET AL: "A novel approach for gene medicine; synthetic poly-L- lysine /serine copolymer enhances bioactivity of antisense oligonucleotides." NUCLEOSIDES AND NUCLEOTIDES, (1997) 16/7-9 (1609-1615). REFS: 10 ISSN: 0732-8311 CODEN: NUNUD5, XP002108844 United States *see in particular abstract and introduction *	1-19
A	MIDOUX P ET AL: "Membrane permeabilization and efficient gene transfer by a peptide containing several histidines." BIOCONJUGATE CHEMISTRY, (1998 MAR-APR) 9 (2) 260-7. JOURNAL CODE: AIT. ISSN: 1043-1802., XP002108845 United States *see in particular the abstract*	1-19

# INTERNATIONAL SEARCH REPORT

International application No.  
PCT/EP 99/08980

## Box I Observations where certain claims were found unsearchable (Continuation of Item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:  
Although claims 11-16 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☒ Claims Nos.:  
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:  
see FURTHER INFORMATION sheet PCT/ISA/210
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.  
☐ No protest accompanied the payment of additional search fees.

## INTERNATIONAL SEARCH REPORT

International Application No. PCT/EP 99/08980

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

### Continuation of Box I.2

The conjugate according to claims 1-8 is not clear in view of the broad and vague definition of both the "monomeric component" and the "protonable residues". The search had to be restricted for economic reasons and was limited to the worked examples and to the general idea underlying the application.  
See Articles 5 and 6 PCT.

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 99/08980

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9822610 A	28-05-1998	FR 2755976 A AU 5123998 A EP 0946744 A	22-05-1998 10-06-1998 06-10-1999
WO 9819710 A	14-05-1998	AU 4873997 A EP 0941123 A	29-05-1998 15-09-1999
US 5627270 A	06-05-1997	US 5693769 A US 5571795 A US 5338837 A AU 687557 B AU 2358295 A CA 2188320 A EP 0756601 A JP 9512270 T NZ 284902 A WO 9529186 A US 5795870 A US 5780444 A AT 166577 T AU 3278593 A BR 9206927 A CA 2117332 A DE 69225719 D DE 69225719 T EP 0618800 A ES 2118932 T HU 70743 A IL 104089 A JP 7503708 T NO 942165 A NZ 246448 A PL 171131 B WO 9311772 A US 5455335 A	02-12-1997 05-11-1996 16-08-1994 26-02-1998 16-11-1995 02-11-1995 05-02-1997 09-12-1997 22-09-1997 02-11-1995 18-08-1998 14-07-1997 15-06-1998 19-07-1993 21-11-1995 24-06-1993 02-07-1998 24-12-1998 12-10-1994 01-10-1998 30-10-1995 09-05-1999 20-04-1995 01-08-1994 28-07-1998 28-03-1997 24-06-1993 03-10-1996
EP 0727223 A	21-08-1996	AU 681192 B AU 7706994 A US 5912300 A CA 2172974 A WO 9509009 A	21-08-1997 18-04-1995 15-06-1999 06-04-1995 06-04-1995